

## A DISCRETE VARIABLE APPROACH FOR INVESTIGATING TUNNELING SPLITTINGS AND VIBRATIONAL WAVE FUNCTIONS IN RARE GAS-ASYMMETRIC TOP HETERODIMERS

EZRA ARUMI ALEXANDER, MARK D. MARSHALL, HELEN O. LEUNG, *Chemistry Department, Amherst College, Amherst, MA, USA.*

A three dimensional discrete variable representation (DVR) is developed for the general case of a rare gas atom interacting with an asymmetric top molecule and applied to the case of argon-haloethylene heterodimers. The position of the rare gas is specified relative to the principal inertial axis system (in a  $I^r$  representation) by the spherical polar coordinates  $(r, \theta, \phi)$ . While the DVR for  $r$  is straightforward, those for  $\theta$  and  $\phi$  presented particular challenges, which are discussed. In common with all DVR approaches, the present DVR is well suited for systems where the intermolecular potential is calculated on a grid of discrete points and provides easy access to the wave functions in the DVR coordinates. The method is applied to the well-characterized argon-*cis*-1,2-difluoroethylene system and used to predict the tunneling splitting and argon-molecule vibrational wave functions in argon-vinyl chloride for which the molecular structure and rotational spectrum remain a puzzle.